

Bi-Para-Mechanical Systems on Lagrangian Distributions

Mehmet Tekkoyun *

Department of Mathematics, Pamukkale University,

20070 Denizli, Turkey

Murat Sari

Department of Mathematics, Pamukkale University,

20070 Denizli, Turkey

January 8, 2009

Abstract

In this work, bi-para-complex analogue of Lagrangian and Hamiltonian systems was introduced on Lagrangian distributions. Yet, the geometric and physical results related to bi-para-dynamical systems were also presented.

Keywords: para-complex geometry, para-Lagrangian, para-Hamiltonian, bi-Lagrangian.

PACS: 02.04; 03.40.

*Corresponding author. E-mail address: tekkoyun@pau.edu.tr; Tel: +902582953616; Fax: +902582953593

1 Introduction

In modern differential geometry, a convenient vector field defined on the tangent and cotangent spaces (or manifolds) which are phase-spaces of velocities and momentum of a given configuration space explains the dynamics of Lagrangians and Hamiltonians. If Q is an m -dimensional configuration manifold and $L(H) : TQ(T^*Q) \rightarrow \mathbf{R}$ is a regular Lagrangian (Hamiltonian) function, then there is a unique vector field ξ (Z_H) on TQ (T^*Q) such that dynamical equations are given by

$$i_\xi \Phi_L = dE_L, \quad i_{Z_H} \Phi_H = dH \quad (1)$$

where Φ_L (Φ_H) indicates the symplectic form. The triples (TQ, Φ_L, ξ) and (T^*Q, Φ_H, Z_H) are called *Lagrangian system* and *Hamiltonian system* on the tangent bundle TQ and on the cotangent bundle T^*Q , respectively.

Mathematical models of mechanical systems are generally described by the Hamiltonian and Lagrangian systems. These models, in particular geometric models in mechanics, are given in some studies [1, 2, 3]. Klein [4] showed that the geometric study of the Lagrangian theory admits an alternative approach without the use of the regular condition on L . It is possible to describe such formalism by using only the differential geometry of the tangent bundles. This is an independent formulation of the Hamiltonian theory. In fact, the role of tangent geometry in Lagrangian theories can be seen to be similar to that of symplectic geometry in Hamiltonian theories [3]. Para-complex analogues of the Euler-Lagrange and Hamiltonian equations were obtained in the framework of Kählerian manifold and the geometric results on a para-complex mechanical systems were found [5]. Symplectic geometry has important roles in some fields of mathematics and physics [6]. As known, Lagrangian foliations on symplectic manifolds are

used in geometric quantization and a connection on a symplectic manifold is an important structure to obtain a deformation quantization. A para-Kählerian manifold M is said to be endowed with an almost bi-para-Lagrangian structure (a bi-para-Lagrangian manifold) if M has two transversal Lagrangian distributions (involutive transversal Lagrangian distributions) D_1 and D_2 [6].

In the above studies; although para-complex geometry and para-complex mechanical systems were analyzed successfully, they have not dealt with bi-para-complex mechanical systems on Lagrangian distributions. In this Letter, therefore, equations related to bi-para-mechanical systems on Lagrangian distributions used in obtaining geometric quantization have been presented.

2 Preliminaries

Throughout this Letter, all mathematical objects and mappings are assumed to be smooth, i.e. infinitely differentiable and Einstein convention of summarizing is adopted, and also $1 \leq i \leq m$. A , $\mathcal{F}(M)$, $\chi(M)$ and $\Lambda^1(M)$ denote the set of para-complex numbers, the set of para-complex functions on M , the set of para-complex vector fields on M and the set of para-complex 1-forms on M , respectively. Since a para-manifold has two transversal Lagrangian distributions; instead of *para* term, *bi-para* term is used for the structures or the mentioned manifold. All structures on the differential manifold M introduced in [6, 7] can be extended in the following subsections.

2.1 Bi-para-complex manifolds

Definition 1: A *product structure* J on a manifold M is a $(1,1)$ tensor field J on M such that $J^2 = I$. The pair (M, J) is called a *product manifold*. A *bi-para-complex manifold* (M, J, D_1, D_2) is a product manifold (M, J) and then one can define a $(1,1)$ tensor field J by $J|_{D_1} = I$ and $J|_{D_2} = -I$. Obviously, $J^2 = I$ and the Nijenhuis $(1,2)$ tensor field N_J vanishes iff D_1 and D_2 distributions are involutive.

Let (x^i, y^i) be a real coordinate system on a neighborhood U of any point p of M , and let $\{(\frac{\partial}{\partial x^i})_p, (\frac{\partial}{\partial y^i})_p\}$ and $\{(dx^i)_p, (dy^i)_p\}$ be natural bases over R of the tangent space $T_p(M)$ and the cotangent space $T_p^*(M)$ of M , respectively. Then the definitions can be given by

$$J(\frac{\partial}{\partial x^i}) = \frac{\partial}{\partial y^i}, \quad J(\frac{\partial}{\partial y^i}) = \frac{\partial}{\partial x^i}. \quad (2)$$

Let $z^i = x^i + \mathbf{j} y^i$, $\mathbf{j}^2 = -1$, also be a para-complex local coordinate system on a neighborhood U of any point p of M . The vector fields can then be shown:

$$(\frac{\partial}{\partial z^i})_p = \frac{1}{2}\{(\frac{\partial}{\partial x^i})_p - \mathbf{j}(\frac{\partial}{\partial y^i})_p\}, \quad (\frac{\partial}{\partial \bar{z}^i})_p = \frac{1}{2}\{(\frac{\partial}{\partial x^i})_p + \mathbf{j}(\frac{\partial}{\partial y^i})_p\}. \quad (3)$$

And the dual covector fields are:

$$(dz^i)_p = (dx^i)_p + \mathbf{j}(dy^i)_p, \quad (d\bar{z}^i)_p = (dx^i)_p - \mathbf{j}(dy^i)_p \quad (4)$$

which represent the bases of the tangent space $T_p(M)$ and cotangent space $T_p^*(M)$ of M , respectively. Then the following expression can be found

$$J(\frac{\partial}{\partial z^i}) = -\mathbf{j}\frac{\partial}{\partial \bar{z}^i}, \quad J(\frac{\partial}{\partial \bar{z}^i}) = \mathbf{j}\frac{\partial}{\partial z^i}. \quad (5)$$

The dual endomorphism J^* of the cotangent space $T_p^*(M)$ at any point p of manifold M satisfies that $J^{*2} = I$, and is defined by

$$J^*(dz^i) = -\mathbf{j}d\bar{z}^i, \quad J^*(d\bar{z}^i) = \mathbf{j}dz^i. \quad (6)$$

2.2 Bi-para-complex modules and bi-para-Hermitian and bi-para-Kählerian manifolds

Here, some definitions and structures on para-Hermitian and para-Kählerian manifold M introduced in [6, 7] are extended to the distributions D_1 and D_2 .

Let V^A be a commutative group $(V, +)$ endowed with a structure of unitary module over the ring A of para-complex numbers. Let V^R denote the group $(V, +)$ endowed with the structure of real vector space inherited from the restriction of scalars to R . The vector space V^R will then be called the real vector space associated to V^A . Setting

$$J(u) = ju, \quad P^+(u) = e^+u, \quad P^-(u) = e^-u, \quad u \in V^A, \quad (7)$$

the expressions

$$\begin{aligned} J^2 &= 1_V, \quad P^{+2} = P^+, \quad P^{-2} = P^-, \quad P^+ \circ P^- = P^- \circ P^+ = 0 \\ P^+ + P^- &= 1_V, \quad P^+ - P^- = J, \\ P^- &= (1/2)(1_V - J), \quad P^+ = (1/2)(1_V + J), \\ j^2 &= 1, \quad e^{+2} = e^+, \quad e^{-2} = e^-, \quad e^+ \circ e^- = e^- \circ e^+ = 0, \\ e^+ + e^- &= 1, \quad e^+ - e^- = j, \\ e^- &= (1/2)(1 - j), \quad e^+ = (1/2)(1 + j). \end{aligned} \quad (8)$$

can be written. Also, it is found that

$$\begin{aligned} J\left(\frac{\partial}{\partial x^i}\right) &= \frac{\partial}{\partial y^i} = \mathbf{j} \frac{\partial}{\partial x^i}, \quad J\left(\frac{\partial}{\partial y^i}\right) = \frac{\partial}{\partial x^i} = \mathbf{j} \frac{\partial}{\partial y^i}, \\ P^\mp\left(\frac{\partial}{\partial z^i}\right) &= -e^\mp \frac{\partial}{\partial z^i}, \quad P^\mp\left(\frac{\partial}{\partial \bar{z}^i}\right) = e^\mp \frac{\partial}{\partial \bar{z}^i}, \\ P^{*\mp}(dz^i) &= -e^\mp dz^i, \quad P^{*\mp}(d\bar{z}^i) = e^\mp d\bar{z}^i. \end{aligned} \quad (9)$$

Definition 2: An *almost bi-para-Hermitian manifold* (M, P^+, P^-, D_1, D_2) is a differentiable manifold M with an almost product structure (P^+, P^-) and a pseudo-Riemannian metric

g , compatible in the sense that

$$g((P^+ - P^-)(X), Y) + g(X, (P^+ - P^-)(Y)) = 0, \quad \forall X, Y \in \chi(D_1) \text{ (or } \chi(D_2)). \quad (10)$$

A *bi-para-Hermitian manifold* is a manifold with an integrable almost bi-para-Hermitian structure (g, P^+, P^-) . Let us take an almost bi-para-Hermitian manifold (M, P^+, P^-, D_1, D_2) . Then, 2-covariant skew tensor field Φ defined by $\Phi(X, Y) = g(X, (P^+ - P^-)(Y))$ is called *fundamental 2-form*. An almost bi-para-Hermitian manifold (M, P^+, P^-, D_1, D_2) is so-called as an *almost bi-para-Kählerian manifold* M if Φ is closed. A bi-para-Hermitian manifold (M, P^+, P^-, D_1, D_2) can be said to be a *bi-para-Kählerian manifold* M if Φ is closed.

3 Bi-Para-Lagrangians

In this section, bi-para-Euler-Lagrange equations and a bi-para-mechanical system can be obtained for classical mechanics structured under the consideration of the basis $\{e^+, e^-\}$ on Lagrangian distributions D_1 and D_2 of bi-para-Kählerian manifold M .

Let (P^+, P^-) be an almost bi-para-complex structure on the bi-para-Kählerian manifold M , and $(z^i, \bar{z}^i)$ be its complex structures. Let semispray be the vector field ξ given by

$$\begin{aligned} \xi &= e^+(\xi^{i+} \frac{\partial}{\partial z^i} + \bar{\xi}^{i+} \frac{\partial}{\partial \bar{z}^i}) + e^-(\xi^{i-} \frac{\partial}{\partial z^i} + \bar{\xi}^{i-} \frac{\partial}{\partial \bar{z}^i}); \\ z^i &= z^{i+} e^+ + z^{i-} e^-; \dot{z}^i = \dot{z}^{i+} e^+ + \dot{z}^{i-} e^- = \xi^{i+} e^+ + \xi^{i-} e^-; \\ \bar{z}^i &= \bar{z}^{i+} e^+ + \bar{z}^{i-} e^-; \dot{\bar{z}}^i = \dot{\bar{z}}^{i+} e^+ + \dot{\bar{z}}^{i-} e^- = \bar{\xi}^{i+} e^+ + \bar{\xi}^{i-} e^-; \end{aligned} \quad (11)$$

where the dot indicates the derivative with respect to time t . The vector field denoted by

$V = (P^+ - P^-)(\xi)$ and given by

$$(P^+ - P^-)(\xi) = e^+(-\xi^{i+} \frac{\partial}{\partial z^i} + \bar{\xi}^{i+} \frac{\partial}{\partial \bar{z}^i}) + e^-(\xi^{i-} \frac{\partial}{\partial z^i} - \bar{\xi}^{i-} \frac{\partial}{\partial \bar{z}^i}) \quad (12)$$

is called *bi-para-Liouville vector field* on the bi-para-Kählerian manifold M . The maps given by $T, P : M \rightarrow A$ such that $T = \frac{1}{2}m_i(\bar{z}^i)^2 = \frac{1}{2}m_i(\dot{z}^i)^2, P = m_i g h$ are called *the kinetic energy* and *the potential energy of the system*, respectively. Here m_i, g and h stand for mass of a mechanical system having m particles, the gravity acceleration and distance to the origin of a mechanical system on the bi-para-Kählerian manifold, respectively. Then $L : M \rightarrow A$ is a map that satisfies the conditions; i) $L = T - P$ is a *bi-para-Lagrangian function*, ii) the function given by $E_L = V(L) - L$ is a *bi-para-energy function*.

The operator $i_{(P^+ - P^-)}$ induced by $P^+ - P^-$ and shown by

$$i_{P^+ - P^-}\omega(Z_1, Z_2, \dots, Z_r) = \sum_{i=1}^r \omega(Z_1, \dots, (P^+ - P^-)Z_i, \dots, Z_r) \quad (13)$$

is said to be *vertical derivation*, where $\omega \in \wedge^r M$, $Z_i \in \chi(M)$. The *vertical differentiation* $d_{(P^+ - P^-)}$ is defined by

$$d_{(P^+ - P^-)} = [i_{(P^+ - P^-)}, d] = i_{(P^+ - P^-)}d - di_{(P^+ - P^-)} \quad (14)$$

where d is the usual exterior derivation. For an almost para-complex structure $P^+ - P^-$, the closed bi-para-Kählerian form is the closed 2-form given by $\Phi_L = -dd_{(P^+ - P^-)}L$ such that

$$d_{(P^+ - P^-)} = e^+B - e^-B : \mathcal{F}(M) \rightarrow \wedge^1 M \quad (15)$$

where

$$B = -\frac{\partial}{\partial z^i}dz^i + \frac{\partial}{\partial \bar{z}^i}d\bar{z}^i.$$

Then

$$\Phi_L = e^+C - e^-C \quad (16)$$

where

$$C = \frac{\partial^2 L}{\partial z^j \partial z^i} dz^j \wedge dz^i - \frac{\partial^2 L}{\partial z^j \partial \bar{z}^i} dz^j \wedge d\bar{z}^i + \frac{\partial^2 L}{\partial \bar{z}^j \partial z^i} d\bar{z}^j \wedge dz^i - \frac{\partial^2 L}{\partial \bar{z}^j \partial \bar{z}^i} d\bar{z}^j \wedge d\bar{z}^i.$$

Let ξ be the second order differential equations given by **Eq.** (1) and

$$\begin{aligned}
i_\xi \Phi_L = \Phi_L(\xi) = & e^+ [\xi^{i+} \frac{\partial^2 L}{\partial z^j \partial z^i} \delta_i^j dz^i - \xi^{i+} \frac{\partial^2 L}{\partial z^j \partial z^i} dz^j - \xi^{i+} \frac{\partial^2 L}{\partial z^j \partial \bar{z}^i} \delta_i^j d\bar{z}^i + \bar{\xi}^{i+} \frac{\partial^2 L}{\partial z^j \partial \bar{z}^i} dz^j \\
& - \xi^{i+} \frac{\partial^2 L}{\partial \bar{z}^j \partial z^i} d\bar{z}^j + \bar{\xi}^{i+} \frac{\partial^2 L}{\partial \bar{z}^j \partial z^i} \delta_i^j dz^i - \bar{\xi}^{i+} \frac{\partial^2 L}{\partial \bar{z}^j \partial \bar{z}^i} \delta_i^j d\bar{z}^i + \bar{\xi}^{i+} \frac{\partial^2 L}{\partial \bar{z}^j \partial \bar{z}^i} d\bar{z}^j] \\
& e^- [-\xi^{i-} \frac{\partial^2 L}{\partial z^j \partial z^i} \delta_i^j dz^i + \xi^{i-} \frac{\partial^2 L}{\partial z^j \partial z^i} dz^j + \xi^{i-} \frac{\partial^2 L}{\partial z^j \partial \bar{z}^i} \delta_i^j d\bar{z}^i - \bar{\xi}^{i-} \frac{\partial^2 L}{\partial z^j \partial \bar{z}^i} dz^j \\
& + \xi^{i-} \frac{\partial^2 L}{\partial \bar{z}^j \partial z^i} d\bar{z}^j - \bar{\xi}^{i-} \frac{\partial^2 L}{\partial \bar{z}^j \partial z^i} \delta_i^j dz^i + \bar{\xi}^{i-} \frac{\partial^2 L}{\partial \bar{z}^j \partial \bar{z}^i} \delta_i^j d\bar{z}^i - \bar{\xi}^{i-} \frac{\partial^2 L}{\partial \bar{z}^j \partial \bar{z}^i} d\bar{z}^j].
\end{aligned} \tag{17}$$

Since the closed bi-para-Kählerian form Φ_L on M is in the structure of bi-para-symplectic, it is found

$$E_L = e^+ (-\xi^{i+} \frac{\partial L}{\partial z^i} + \bar{\xi}^{i+} \frac{\partial L}{\partial \bar{z}^i}) + e^- (\xi^{i-} \frac{\partial L}{\partial z^i} - \bar{\xi}^{i-} \frac{\partial L}{\partial \bar{z}^i}) - L \tag{18}$$

and thus

$$\begin{aligned}
dE_L = & -\xi^{i+} \frac{\partial^2 L}{\partial z^j \partial z^i} dz^j e^+ + \bar{\xi}^{i+} \frac{\partial^2 L}{\partial z^j \partial \bar{z}^i} dz^j e^+ - \xi^{i+} \frac{\partial^2 L}{\partial \bar{z}^j \partial z^i} d\bar{z}^j e^+ + \bar{\xi}^{i+} \frac{\partial^2 L}{\partial \bar{z}^j \partial \bar{z}^i} d\bar{z}^j e^+ \\
& - \frac{\partial^2 L}{\partial z^j} dz^i + \xi^{i-} \frac{\partial^2 L}{\partial z^j \partial z^i} dz^j e^- - \bar{\xi}^{i-} \frac{\partial^2 L}{\partial z^j \partial \bar{z}^i} dz^j e^- + \xi^{i-} \frac{\partial^2 L}{\partial \bar{z}^j \partial z^i} d\bar{z}^j e^- \\
& - \bar{\xi}^{i-} \frac{\partial^2 L}{\partial \bar{z}^j \partial \bar{z}^i} d\bar{z}^j e^- - \frac{\partial^2 L}{\partial \bar{z}^j} d\bar{z}^i.
\end{aligned} \tag{19}$$

With the use of **Eq.** (1), the following expression can be obtained:

$$\begin{aligned}
& \xi^{i+} \frac{\partial^2 L}{\partial z^j \partial z^i} dz^j e^+ - \xi^{i+} \frac{\partial^2 L}{\partial z^j \partial \bar{z}^i} d\bar{z}^j e^+ + \bar{\xi}^{i+} \frac{\partial^2 L}{\partial \bar{z}^j \partial z^i} dz^j e^+ - \bar{\xi}^{i+} \frac{\partial^2 L}{\partial \bar{z}^j \partial \bar{z}^i} d\bar{z}^j e^+ \\
& - \xi^{i-} \frac{\partial^2 L}{\partial z^j \partial z^i} dz^j e^- + \xi^{i-} \frac{\partial^2 L}{\partial z^j \partial \bar{z}^i} d\bar{z}^j e^- - \bar{\xi}^{i-} \frac{\partial^2 L}{\partial \bar{z}^j \partial z^i} dz^j e^- + \bar{\xi}^{i-} \frac{\partial^2 L}{\partial \bar{z}^j \partial \bar{z}^i} d\bar{z}^j e^- + \frac{\partial^2 L}{\partial z^j} dz^i + \frac{\partial^2 L}{\partial \bar{z}^j} d\bar{z}^i = 0.
\end{aligned} \tag{20}$$

If a curve denoted by $\alpha : A \rightarrow M$ is considered to be an integral curve of ξ , then the equations given in the following are

$$\begin{aligned}
& \left[\xi^{i+} \frac{\partial^2 L}{\partial z^j \partial z^i} + \bar{\xi}^{i+} \frac{\partial^2 L}{\partial \bar{z}^j \partial z^i} \right] dz^j e^+ - \left[\xi^{i-} \frac{\partial^2 L}{\partial z^j \partial z^i} + \bar{\xi}^{i-} \frac{\partial^2 L}{\partial \bar{z}^j \partial z^i} \right] dz^j e^- \\
& - \left[\xi^{i+} \frac{\partial^2 L}{\partial z^j \partial \bar{z}^i} + \bar{\xi}^{i+} \frac{\partial^2 L}{\partial \bar{z}^j \partial \bar{z}^i} \right] d\bar{z}^j e^+ + \left[\xi^{i-} \frac{\partial^2 L}{\partial z^j \partial \bar{z}^i} + \bar{\xi}^{i-} \frac{\partial^2 L}{\partial \bar{z}^j \partial \bar{z}^i} \right] d\bar{z}^j e^- + \frac{\partial L}{\partial z^j} dz^i + \frac{\partial L}{\partial \bar{z}^j} d\bar{z}^i = 0
\end{aligned} \tag{21}$$

or alternatively

$$(e^+ - e^-) \left(\left[\xi^{i+} \frac{\partial}{\partial z^{j^i}} + \bar{\xi}^{i+} \frac{\partial}{\partial \bar{z}^j} \right] e^+ + \left[\xi^{i-} \frac{\partial}{\partial z^j} + \bar{\xi}^{i-} \frac{\partial}{\partial \bar{z}^j} \right] e^- \right) \left(\frac{\partial L}{\partial z^i} \right) dz^j$$

$$(e^+ - e^-) \left(- \left[\xi^{i+} \frac{\partial}{\partial z^{j^i}} + \bar{\xi}^{i+} \frac{\partial}{\partial \bar{z}^j} \right] e^+ - \left[\xi^{i-} \frac{\partial}{\partial z^j} + \bar{\xi}^{i-} \frac{\partial}{\partial \bar{z}^j} \right] e^- \right) \left(\frac{\partial L}{\partial \bar{z}^i} \right) d\bar{z}^j + \frac{\partial L}{\partial z^j} dz^j + \frac{\partial L}{\partial \bar{z}^j} d\bar{z}^j = 0.$$

Then the following equations are found:

$$(e^+ - e^-) \frac{\partial}{\partial t} \left(\frac{\partial L}{\partial z^i} \right) + \frac{\partial L}{\partial z^i} = 0, \quad -(e^+ - e^-) \frac{\partial}{\partial t} \left(\frac{\partial L}{\partial \bar{z}^i} \right) + \frac{\partial L}{\partial \bar{z}^i} = 0 \quad (22)$$

or

$$(e^+ - e^-) \frac{\partial}{\partial t} \left(\frac{\partial L}{\partial z^i} \right) + \frac{\partial L}{\partial z^i} = 0, \quad (e^+ - e^-) \frac{\partial}{\partial t} \left(\frac{\partial L}{\partial \bar{z}^i} \right) - \frac{\partial L}{\partial \bar{z}^i} = 0. \quad (23)$$

Thus the equations obtained in **Eq.** (23) are seen to be a *bi-para-Euler-Lagrange equations* on the distributions D_1 and D_2 , and then the triple (M, Φ_L, ξ) is seen to be a *bi-para-mechanical system* with taking into account the basis $\{e^+, e^-\}$ on the distributions D_1 and D_2 .

4 Bi-Para-Hamiltonians

Here, bi-para-Hamiltonian equations and bi-para-Hamiltonian mechanical system for classical mechanics structured on the distributions D_1 and D_2 are derived.

Let M be a bi-para-Kählerian manifold and $\{z_i, \bar{z}_i\}$ be its complex coordinates. Let $\{\frac{\partial}{\partial z_i}|_p, \frac{\partial}{\partial \bar{z}_i}|_p\}$ and $\{dz_i|_p, (d\bar{z}_i)_p\}$, also, be bases over para-complex number A of tangent space $T_p(M)$ and cotangent space $T_p^*(M)$ of M , respectively. Let us assume that an almost bi-para-complex structure, a bi-para-Liouville form and a bi-para-complex 1-form on the distributions D_1 and D_2 are shown by $P^{*+} - P^{*-}$, λ and ω , respectively. Then $\omega = \frac{1}{2}[(z_i d\bar{z}_i + \bar{z}_i dz_i)e^+ + (z_i d\bar{z}_i + \bar{z}_i dz_i)e^-]$ and $\lambda = (P^{*+} - P^{*-})(\omega) = \frac{1}{2}[(z_i d\bar{z}_i - \bar{z}_i dz_i)e^+ - (z_i d\bar{z}_i - \bar{z}_i dz_i)e^-]$. It is concluded that if Φ is a closed bi-para-Kählerian form on the distributions D_1 and D_2 , then Φ is also a bi-para-symplectic structure on D_1 and D_2 .

Consider that bi-para-Hamiltonian vector field Z_H associated with bi-para-Hamiltonian energy H is given by

$$Z_H = (Z_i \frac{\partial}{\partial z_i} + \bar{Z}_i \frac{\partial}{\partial \bar{z}_i})e^+ + (Z_i \frac{\partial}{\partial z_i} + \bar{Z}_i \frac{\partial}{\partial \bar{z}_i})e^-. \quad (24)$$

Then

$$\Phi = -d\lambda = (e^+ - e^-)(d\bar{z}_i \wedge dz_i)$$

and

$$i_{Z_H} \Phi = \Phi(Z_H) = (\bar{Z}_i dz_i - Z_i d\bar{z}_i)e^+ + (-\bar{Z}_i dz_i + Z_i d\bar{z}_i)e^-. \quad (25)$$

Moreover, the differential of bi-para-Hamiltonian energy is obtained as follows:

$$dH = (\frac{\partial H}{\partial z_i} dz_i + \frac{\partial H}{\partial \bar{z}_i} d\bar{z}_i)e^+ + (\frac{\partial H}{\partial z_i} dz_i + \frac{\partial H}{\partial \bar{z}_i} d\bar{z}_i)e^-. \quad (26)$$

By means of **Eq.**(1), using **Eq.** (25) and **Eq.** (26), the bi-para-Hamiltonian vector field is found to be

$$Z_H = (-\frac{\partial H}{\partial \bar{z}_i} \frac{\partial}{\partial z_i} + \frac{\partial H}{\partial z_i} \frac{\partial}{\partial \bar{z}_i})e^+ + (\frac{\partial H}{\partial \bar{z}_i} \frac{\partial}{\partial z_i} - \frac{\partial H}{\partial z_i} \frac{\partial}{\partial \bar{z}_i})e^-. \quad (27)$$

Suppose that a curve

$$\alpha : I \subset A \rightarrow M \quad (28)$$

be an integral curve of the bi-para-Hamiltonian vector field Z_H , i.e.,

$$Z_H(\alpha(t)) = \dot{\alpha}, \quad t \in I. \quad (29)$$

In the local coordinates, it is obtained that

$$\alpha(t) = (z_i(t), \bar{z}_i(t)) \quad (30)$$

and

$$\dot{\alpha}(t) = (\frac{dz_i}{dt} \frac{\partial}{\partial z_i} + \frac{d\bar{z}_i}{dt} \frac{\partial}{\partial \bar{z}_i})e^+ + (\frac{dz_i}{dt} \frac{\partial}{\partial z_i} + \frac{d\bar{z}_i}{dt} \frac{\partial}{\partial \bar{z}_i})e^-. \quad (31)$$

Consideration of **Eqs.** (29), (27), (31), the following results can be obtained

$$\frac{dz_i}{dt}e^+ = -\frac{\partial H}{\partial \bar{z}_i}e^+, \frac{dz_i}{dt}e^- = \frac{\partial H}{\partial \bar{z}_i}e^-, \frac{d\bar{z}_i}{dt}e^+ = \frac{\partial H}{\partial z_i}e^+, \frac{d\bar{z}_i}{dt}e^- = -\frac{\partial H}{\partial z_i}e^- \quad (32)$$

or in a more simplified form:

$$\frac{dz_i}{dt} = -(e^+ - e^-)\frac{\partial H}{\partial \bar{z}_i}, \quad \frac{d\bar{z}_i}{dt} = (e^+ - e^-)\frac{\partial H}{\partial z_i}. \quad (33)$$

Hence, the equations obtained in **Eq.** (33) are seen to be *bi-para Hamiltonian equations* on the distributions D_1 and D_2 , and then the triple (M, Φ_L, Z_H) is seen to be a *bi-para-Hamiltonian mechanical system* with the use of basis $\{e^+, e^-\}$ on the distributions D_1 and D_2 .

5 Conclusion

This study has shown to exist physical proof of the mathematical equality given by $M = D_1 \oplus D_2$. Also, formalisms of Lagrangian and Hamiltonian mechanics have intrinsically been described with taking into account the basis $\{e^+, e^-\}$ on Lagrangian distributions D_1 and D_2 of bi-para-Kählerian manifold M .

Bi-para-Lagrangian and bi-para-Hamiltonian models arise to be a very important tool since they present a simple method to describe the model for bi-para-mechanical systems. In solving problems in classical mechanics, the bi-para-complex mechanical system will then be easily usable model. With the use of the corresponding approach, thus, a differential equation resulted in mechanics is seen to have a non-trivial solution.

Since physical phenomena, as well-known, do not take place all over the space, a new model for dynamical systems on subspaces is needed. Therefore, equations (23) and (33) are only

considered to be a first step to realize how bi-para-complex geometry has been used in solving problems in different physical area.

For further research, bi-para-complex Lagrangian and Hamiltonian vector fields derived here are suggested to deal with problems in electrical, magnetical and gravitational fields of physics.

References

- [1] Crampin M. On the differential geometry of the Euler-Lagrange equations, and the inverse problem of Lagrangian dynamics. J. Phys. A: Math. Gen, 1981; 14: 2567-2575
- [2] Nutku N. Hamiltonian Formulation of KdV Equation. J. Math. Phys. 1984; 25: 2007-2008
- [3] De Leon M. and Rodrigues PR. Methods of Differential Geometry in Analytical Mechanics. North-Holland Mathematics Studies 152; Elsevier; Amsterdam; 1989.
- [4] Klein J. Espaces variationnels et Mécanique. Ann. Inst. Fourier (Grenoble). 1962; 12: 1-124
- [5] Tekkoyun M. On Para Euler-Lagrange and Para-Hamiltonian equations. Phys. Lett. A, 2005; 340: 7-11
- [6] Etayo F. and Santanarúa R. and Trías UR. The geometry of a biLagrangian manifold. Diff. Geom. and its Appl. 2006; 24: 33-59
- [7] V. Cruceanu, P.M. Gadea, J. M. Masqué, Para-Hermitian and Para- Kähler Manifolds, Supported by the commission of the European Communities' Action for Cooperation in Sciences and Technology with Central Eastern European Countries n. ERB3510PL920841.